


Wayne County Center

Livestock News

Sep & Oct 2020



Inside This Issue

- 1 Important Information
- 2 Animal Waste Mgmt.
- 3 Don't Ignore Hybrid Vigor
- 4 More Than Just Ryegrass
- 5 Managing Internal Parasites in Sheep and Goats
- 6 Biosecurity During COVID-19 for Horse Owners
- 7 Virtual Livestock Options for NC Youth
- 8 Water Quality

Contact Us

NC Cooperative Extension
Cumberland County Center
3114 B Wayne Memorial Dr
Goldsboro, NC 27534
Wayne.ces.ncsu.edu

Stefani Sykes
Extension Agent, Livestock
Stefani.sykes@waynegov.com
919-731-1525

NC State University and N.C. A&T State University commit themselves to positive action to secure equal opportunity and prohibit discrimination and harassment regardless of age, color, disability, family and marital status, genetic information, national origin, political beliefs, race, religion, sexual identity (including pregnancy) and veteran status. NC State, N.C. A&T, U.S. Department of Agriculture, and local governments cooperating.

NC State Extension works in tandem with N.C. A&T State University, as well as federal, state and local governments, to form a strategic partnership known as N.C. Cooperative Extension.

Fall Virtual Beef Cattle Series

Join us online for this three-part webinar series to learn about feed supplementation based on hay quality, managing first-calf heifers, and freezer beef 101. This series will take place on October 15, November 19, and December 17 from 6:30-7:30 PM. You can choose to register for one, two, or all three sessions. These webinars are FREE!

The webinars will be delivered through Zoom, which is an online video communications platform that is free for users. You can download the free app on your phone or you can join by computer. If you cannot join by smartphone or computer, you can call in and listen from any phone. To register, go to the following link: <https://go.ncsu.edu/fallbeefseries>

Upcoming Pesticide Classes

- Z Training (fumigation) :** Sept. 23
8:00-9:00 am
 - V Training:** Sept. 23
9:30-11:30 am
 - X Training:** Sep 23
1:00-3:00 pm
- Call 919-731-1525 to register!**

Coronavirus Food Assistance Program (CFAP)

CFAP provides financial assistance to producers of agricultural commodities who have suffered a five-percent or greater price decline or who had losses due to market supply chain disruptions due to COVID-19. Eligible commodities include:

- Non-specialty crops (malting barley, canola, corn, upland cotton, millet, oats, soybeans, sorghum, sunflowers, durum wheat, and hard red spring wheat)
 - Wool
- Livestock (cattle, hogs, and sheep - lambs and yearlings)
 - Dairy
- Specialty crops (fruits, vegetables, nuts, beans, and mushrooms)

Applications are being accepted now until September 11, 2020. Producers should apply through their local Farm Service Agency office. More information about the program can be found at farmers.gov/cfap.

Call 919-734-5281 ext #2 for more information.

Hay Directory

North Carolina Department of Agriculture's Hay Alert is at <http://www.ncagr.gov/HayAlert/>. It lists people selling hay or looking for hay to buy. It is free to list your hay.

For any meeting listed, persons with disabilities may request accommodations to participate by contacting the Extension Office where the meeting will be held by phone, email, or in person at least 7 days prior to the event.

Disclaimer - The use of brand names and any mention or listing of commercial products or services in this publication does not imply endorsement by North Carolina State University nor discrimination against similar products or services not mentioned.

Fall Recertification Classes for Animal Waste Operators

By: Eve Honeycutt, Livestock Extension Agent with N.C. Cooperative Extension in Lenoir and Greene Counties

This year, Extension is planning to host animal waste recertification classes in a variety of ways due to COVID-19. Currently, the Governor's order says we can have no more than 10 people gathered inside. With that in mind, we have two options for those that need their credit hours before December 31, 2020:

- Attend a small face-to-face class offered by Extension (9 or less participants and one Extension Agent)
- Attend two (3 hour) statewide Zoom session (online)

We are asking that those of you who do not HAVE to have your credit hours by December 31, 2020, please attend the online training or wait until another year.

The dates and topics for the online classes are below. Each session is worth 3 hours CEC:

November 10, 5-8pm: <https://november10cec.eventbrite.com>

- Best Management Practices for Sampling Manures (30min) - Mahmoud Sharara, NCSU
- Weather- Hurricanes, Tornadoes, Flooding- (1 hour)- National Weather Service
- Hay Production- (30min) Dennis Hancock, formerly UGA
- Proper Tissue Sampling- (30min)- Steph Kulesza, NCSU

- Nutrient Mangement Planning and Cropping Systems- (30min)- Steph Kulesza, NCSU

December 3, 9am-12pm: <https://december3cec.eventbrite.com>

- Sludge Management and Utilization Opportunities-(1 hour) Mahmoud Sharara, NCSU
- NC Pork Council- Legislative Updates (1 hour)- Angie Maier, NCPC
- What to expect on a DEQ inspection (30min)- Megan Stille, Washington Regional Office
- African Swine Fever update (30min)- Jon Holt, NCSU

December 16, 1-4pm: <https://december16cec.eventbrite.com>

- Using Animal Waste in Forestry- (30min)- Colby Lambert, NCSU Extension
- Bermudagrass Fertilization and Issues- (1 hour)- Georgia Love, NCDA Regional Agronomist
- Crop Production- (30min)- NCSU Crop Science Department
- Research on Identifying DNA of Waste Products- (1 hour)- NC Pork Council cooperation

Registration is open for this events. Use the link provided above to register for the class. After you register, a zoom link will be sent via email a few days before the class.

Challenging Times for the Pork Industry

By: Eve Honeycutt, Livestock Extension Agent with N.C. Cooperative Extension in Lenoir and Greene Counties

As you know, the pork industry has also been affected by the events of 2020. There are several reasons for the challenges facing the industry right now.

- Not as many people eating pork- the slow down of the food service industry (due to COVID-19) pushed demand for pork products way down.
- Since 2014, the total number of hogs and pigs have increased by 20 million head. This is due to increased productivity in the industry.
- Because of the increase in numbers, we have more pigs than we can physically process. We increased numbers but did not increase slaughter capacity.
- Even with the new trade deal with China, they have not met the target for phase one of the deal.

The result of this increased supply and decreased demand, is extremely low prices. This creates a serious and long term decrease in returns for the industry overall. Since the United States is a low cost producer of pork, it makes sense that this supply and demand curve will balance back out, but that timeframe is unknown.

Most of the companies that own sows have decreased breeding considerably. Some companies have had to make exceptionally hard decisions for the future of their business. This has created a slow down or complete shut-down of supply for contract growers. Understandably, growers affected are very concerned for their livelihood and the future of their farm.

Once the contract runs out, the growers who find themselves without hogs have to consider a few options.

Patience- New opportunities will become available in

the industry. The ability for a grower to wait for an upturn is a very personal decision, and can be affected by the amount of debt owed.

Stand empty- If you choose to get out of the industry, you must remember that the lagoon and spray-fields must still be maintained as stated in the permit. As long as the farm has a lagoon (and permit), it still has the value associated with a hog farm in NC. A farm that stands empty for less than 5 years can re-populate with no additional requirements. This 5 year clock can be reset by populating the farm with at least 10% of the steady-state live weight or 250 hogs, whichever is fewer, for a period of at least 45 days. Farms empty for more than 5 years have considerable requirements to meet before repopulating. Farms empty for more than 10 years can not repopulate.

Close farm- Closing the farm means also closing the lagoon and rescinding the permit. This means confined hogs (over 250 head) can never be placed back on this site. Closing the lagoon also involves quite a bit of planning and covering the expenses associated with the closure.

Independent production- There are niche markets that are interested in buying more pork. However, being an independent producer can be quite challenging. If you are interested in discovering more about producing pigs independently, we have a team of specialists and economists at NCSU that can assist you by providing information to help you make an informed decision.

Please let us know if Extension can help you in any way. We can find answers to your questions at any time.

Don't Ignore Hybrid Vigor

By: Paul Gonzalez, Livestock Extension Agent with N.C. Cooperative Extension in Sampson County

Many producers have gotten away from planned cross-breeding systems. I see this as an issue everywhere I go, not just in Sampson County. It is due partly to the ready availability and ease with which an Angus bull can be purchased and partly to the ease of management when using only one sire breed; but, is mainly due to the market demand for black cattle. I know some people will not like what I have to say, and I will come across looking like I am Angus-bashing! While that is not my intent and I am not anti-Angus, it will be the main focus of this article due to the fact that Angus is at the center of this dilemma. Keep in mind; however, that what is said here applies to any breed of cattle, not just Angus.

I will start by saying that more and more producers are, whether knowingly or unknowingly, moving toward cow herds that are basically purebred Angus. Year after year, Angus bulls are turned in with the cows; and heifers from those bulls are kept for replacements. If replacements aren't retained, Angus sired heifers or cows are usually purchased to enter the herd. These females are then bred to an Angus bull and the pattern is repeated. Assuming I started with a different breed initially, after using Angus bulls and keeping heifers for four years, I have some heifers now that are 93.75% Angus. This percentage is considered to be purebred by breed associations that allow breeding up. As you can see, in a few more years as you cull older cows and replace them with heifers, you have a herd that is considered purebred Angus. Again, keep in mind the same will be true if you use any breed in the manner described above.

I stated in the opening paragraph that the main reason for this is the market demand for black cattle. The easiest way to assure yourself of getting black cattle is to use an Angus bull. There is very little chance that a black Angus bull carries the red gene anymore so you don't have that concern like you would using a black bull of another breed. However, I would also like to point out that there are homozygous black bulls available in other breeds these days. Another reason is the ease of management. You only have to have one breeding pasture. You don't need more than one bull. And you don't have to keep up with parentage on heifers or how long you have been using a bull if you never switch breeds. A simple rotation of bull breeds every four years will make a big difference in your herd while still keeping management fairly simple. Or you can make life even simpler by using a composite bull. Choose a composite breed and stick with it. You won't get the same amount of heterosis as an initial cross but will stabilize around 67% of the potential gain.

Now to the point of the article. By the continued use of the same sire breed, you are losing the effects of heterosis and giving up what are essentially free pounds. Heterosis, or hybrid vigor, is the improvement shown by crossbred animals over straight bred animals. How much improvement varies depending on the breeds used but for weaning weight will average about 4% for crossbred calves over straight bred calves that are both nursing

straight bred cows. This means you would get another 20 pounds on each 500 lb calf just for switching bulls. If you have twenty-five cows, it is like selling another calf. Not only do the calves weigh more but you also get added survivability in the crossbred calves so you get another 3% heterosis advantage in weaning percent. Let's look at an example.

Say breed A calves average 480 lbs at weaning and breed B calves average 520 lbs. Calves sired by breed A out of breed B cows have weaning weights of 540 lbs and calves by breed B sires out of breed A cows average 520 lbs. The amount of heterosis from the crossbreeding would be figured by subtracting the straight bred average $(480+520)/2=500$ from the crossbred average $(540+520)/2=530$ and dividing that amount, 30 in this example, by the straight bred average and multiplying by 100 $[(30/500)*100=6]$ which yields a 6% heterosis value.

The advantages become even greater if you use a third breed on crossbred cows. You get the added pounds from the calf heterosis in the example above. You also get greater weaning percentage from the crossbred cows, due to higher conception rates, and even greater weaning weights due to increased milk production. In a study conducted at the Fort Robinson Research Station (Cundiff and Gregory, 1977; Gregory and Cundiff, 1980), crossbred cows raising crossbred calves weaned 23.3 percent more calf weight per cow exposed than straightbred cows raising straightbred calves. Two thirds of the advantage was attributed to the maternal heterosis of the cow and one third to the individual heterosis of the calf. Other studies have show increases of up to 28%. Experiments using Brahman/European crosses have demonstrated even greater total increases over the straightbred parents.

It has been proven through research that hybrid vigor will add pounds to your calf crop. In high market price times, it may not seem as significant but when prices are low it is extremely important. This additional weight should not be dismissed to simply chase black hides. Again, let me emphasize that I am not in any way bashing, degrading, or opposing the use of Angus cattle. Angus cattle have made great contributions to the beef industry and absolutely have their place in a well planned and implemented crossbreeding system. It seems though that some producers see them as a silver bullet and have fallen into a straight breeding rut. As I stated earlier, homozygous black cattle are becoming available in all breeds. There are breeders of black Herefords with bulls on the market. And, I have even seen some registered black Charolais! So, you can still meet the demand for black calves using a second breed of bull. One thing I should mention here is to choose breeds which are similar and complement each other. I won't make breed recommendations here but if you would like to discuss, or debate, breed choices and breeding systems, feel free to give me a call.

More Than Just Ryegrass

By: Kelly McCaskill, Livestock Extension Agent with N.C. Cooperative Extension in Moore County

Annual forages are not for every farm, but there are lots of folks that could benefit from including annuals in their pasture mix, even just for a season. Most people know about the use of ryegrass for overseeding a Bermuda pasture but did you know there is a whole world of cool season annual forages out there just waiting to be put to work on your property? Cool season annual forages go way beyond just ryegrass!

Cool season annual forages can be broken down into three main categories: grasses, legumes and brassicas. Depending on your production goals and management style they can be planted individually or as a mix.

Grasses

Cereal Rye-the most common small grain planted as a forage and for good reason. It is pretty forgiving of soil and weather conditions, establishes quickly and produces tons of biomass. This forage does mature quickly and loses quality as it matures so grazing is very timely.

Oats-this forage can be early fall or late winter planted but is most productive in the spring. The broad leaves provide excellent yield and quality. Just like cereal rye, this forage does lose quality as it matures, but not quite as quickly. Oats establish very quickly, usually being ready to graze in less than two months, making it a great 'emergency forage'.

Triticale-this is a hybrid forage, boasting the high yielding properties of wheat and the cold tolerance of rye, making this a well-rounded cool season option.

Legumes

Clover-there are several types of clover that can be planted in combination with other forages to offer high quality grazing, while simultaneously improving your soil. Clover, as well as all other legumes, are nitrogen-fixers, taking nitrogen from the atmosphere and adding it back into the soil. Crimson, Arrowleaf and Ladino clovers are common planted varieties that provide a palatable, nutritious addition to grass systems. Clovers must be allowed ample time to establish before they can be grazed, so they are usually used for spring grazing following a fall seeding.

Peas-there are many pea varieties on the market that are great for grazing. Austrian winter pea, cow-pea, and good ol' field peas are just a few that make excellent forages when in combination with grasses. Peas need to be seeded a little deeper than their

smaller seeded clover cousins but they can still be used in a no-till system, just pay attention to the drill depth.

Vetch-this forage is one that will send a row crop farmer into fits because it has no place in a tobacco or corn field but makes wonderful animal food. Vetch should be planted in combination with other forages such as grasses. Hairy vetch is the most common variety but there are others available that make good forage as well. Vetch is very palatable and has the highest nitrogen fixation potential of all of the legumes, but is also one of the latest forages to come on, usually not being grazable until mid to late spring.

Brassicas

Turnips-these forages are very quick to establish and make a great addition to a mixed forage system. They establish quickly and have the potential to produce tons of biomass. They have a tuber under the ground, that will vary in size depending on the variety, as well as large delicious leaves. All brassicas are high in protein and low in fiber so care should be taken not to overfeed to livestock, which can cause health issues. This can be avoided by using brassicas in a mix of other forages such as grasses and legumes, which are higher in fiber and lower in protein.

Radishes-similar to turnips, this forage provides both above ground foliage as well as a tuber beneath the earth, varying in size depending on the variety. Some varieties of radish boast "tillage" properties due to the size of the tuber. These varieties are ideal for soil with compaction issues.

Kale-this brassica typically does not have a notable underground tuber but does produce a large amount of leaf matter above the ground. It is nutritious, delicious and typically very winter-hardy. Just like the other brassicas, this is one that should be grazed in combination with grasses or a grass/legume mix.

These forages are just the tip of the iceberg as to what is available for annual winter grazing options. Contact your local extension agent to discuss what works best in a mix as well as what dates, depths and rates to use for seeding.

Managing Internal Parasites in Sheep and Goats

By: Anthony Growe, Livestock and Row Crops Extension Agent with N.C. Cooperative Extension in Richmond County

As we near the end of the summer, small ruminant owners should be thinking about internal parasite management. Generally speaking, stomach worm populations, such as the Barberpole worm, accumulate over the summer grazing season. This buildup creates internal parasite issues in the late summer months, especially if there is adequate moisture. Based on recent weather conditions, I suspect we will see heavy parasite loads in our pastures. Most stomach worms like a warm, moist environment. Temperatures around 85 degrees Fahrenheit coupled with high humidity and moisture make an ideal environment for the eggs to hatch and develop into larvae. Since the Barberpole worm is one of the most significant internal parasites in small ruminants, this article will concentrate on managing this particular species.

Before we begin trying to manage internal parasites, it's important to understand their general lifecycle. In our area, the winters are not cold enough to kill all worm larvae so stomach worms can overwinter in the ground or even in the animal's digestive system. The worms that were able to overwinter deep in the stomach of the animal are called arrested larva which lay dormant until they are triggered by a stress such as kidding or lambing. After breaking dormancy, these arrested larvae mature into stomach worms and produce eggs. These eggs are shed from the animal and deposited to the ground in their feces.

About a week or so after worm larvae hatch from their eggs in a manure pellet, they climb up 2 to 3 inches onto grass blades that goats and sheep consume. When the animal ingests the larvae, they attach and feed on the true stomach and intestine linings which can be damaged or irritated. This damage reduces the amount of nutrients that an animal can absorb during digestion. Additionally, some eggs are re-deposited back to the ground in manure, continuing the infectious cycle.

If left unchecked, stomach worms will cause weight loss, stunted growth, anemia, poor conception rates and can even cause death. To get an idea of how heavy a worm load is, periodically examine your animals. Things that usually indicate heavy worm infestations are: thin body condition, weight loss, rough hair coat, pale or white tongue and inner eyelids, and diarrhea. If you are properly feeding your animals but they exhibit these symptoms, then you may need to

implement some management options. Here are some tips goat and sheep owners can use to manage worm loads in their livestock:

- Provide your animals with a nutritious diet for a good immune system
- Divide up large pastures to allow for rotational grazing
- Graze above the worms by keeping pasture grass at least 4 inches high
- Reduce contamination by keeping feed elevated and off the ground
- Keeping water troughs and feed bunks clean will also reduce worm ingestion
- Monitor parasite loads by using FAMACHA or fecal egg counts
- Time dewormings around kidding and lambing season
- Refrain from using worm blocks (it's hard to measure a single animal's intake)
- Feeding copper boluses to goats may help reduce Barber Pole worm infestations
- When using recommended dewormers rotate chemical classes and selectively treat animals to minimize resistance
- Work with a veterinarian who is familiar with internal parasite issues in your area to provide "extra-label" dewormer recommendations.

Internal parasites are a significant issue in small ruminant management. Adopting practices, such as rotational grazing and good sanitation, will help reduce infective parasites in your herds or flock. It is also important that producers employ these best management practices to slow internal parasite resistance to our dewormers so that they can continue to be effective on our target parasites. If you have any questions concerning livestock management, please call your local livestock Extension agent.

Biosecurity During COVID-19 for Horse Owners

By: Taylor Chavis, Livestock Extension Agent with N.C. Cooperative Extension in Robeson County

The horse industry in all fashions have taken a hit because of Covid-19. Many horse shows cancelled, trail locations closed, and boarding facilities under stricter regulations have horse owners “biting at the bit” (pun intended) to get back in the saddle. However, horse owners should take precautions to protect themselves and their horses. Below are some tips for practicing biosecurity during a pandemic at either horse shows, trail riding locations, barns, etc.

According to the CDC, there has been no report of horses being infected with the SARS-coV-2, virus that causes Covid-19. However, biosecurity measures should be in place to prevent the spread of other potential equine diseases.

- Horses should not be co-mingled. One carrier animal can potentially infect not only the other horses at the event, but those pathogens can be taken home and shared with the rest of the horses at the home farm if precautions are not taken.
- Be sure that the trailer used to haul the horses is clean, as well as equipment and tack.
- Talk to your veterinarian about any concerns of horse health, make sure the vaccinations are up to date, discuss any disease concerns in the area (s) to be travelled through, and any other health concerns.
- Don't share tack, grooming supplies, feed pans, or water buckets. Most equine diseases are spread by direct contact. Direct contact not only includes nose-to-nose contact, but contact with surfaces that may have gotten saliva, respiratory secretions, or manure contamination from an infected horse.
- Keep the area clean around your horses. Fully clean and disinfect any stalls to be used before putting your horses in. Do this even if the event host has already promised that the stalls were cleaned and disinfected.
- Keep an eye on your horses for any sign of illness. Check for fever, and keep track of feed and water intake while away from home.
- When returning from an equine event, isolate the horse that has traveled for at least a week. Monitor for any symptoms, like fever, off-feed, etc. If symptoms are seen, veterinary care can be given.
- Clean and disinfect the trailer, tack, and equipment when returning from events.
- Keep your horse's stress level as low as possible. Stress can cause the horse's immune system to become compromised. Continue a familiar feeding regimen to keep your

horse on schedule to avoid stress and colic. If your horse is not used to being in a stall and gets stressed when they are in one, taking them out frequently and hand-walking them can help decrease their stress level.

The CDC recommends these tips below for persons organizing livestock shows:

- Split classes to limit the number of animals in the show ring.
- Limit the number of people and/or animals in a ring to ensure that at least 6 feet is kept between an animal and its handler and other people and animals including judges. The number of exhibitors and animals that can safely fit in a show ring or exhibit area may limit the number of animals that can be shown at one time.
- Provide separate entry and exit points for visitors so they do not need to pass close together while coming into and going out of the ring.
- Stagger activities in washing and grooming areas, or other shared spaces, so that animals from different farms or households do not interact unnecessarily.
- Stagger animal move-in/move-out times to reduce contact between people and animals.
- Consider a “show-and-go” with animals stalled at a trailer and leaving immediately after the show if this will not compromise animal welfare, for example, because of weather conditions such as high temperature and humidity.

Lastly, horse owners should take precautions to protect not only their horses, but themselves. The CDC recommends that persons should:

- Wash their hands frequently for at least 20 seconds with soap and water
- Use a hand sanitizer that has at least 60% alcohol
- Avoid touching eyes, nose, and mouth with unwashed hands
- Avoid close contact with other persons
- Maintain at least 6 feet
- Cover your nose and mouth with a mask
- Clean and disinfect frequently touched surfaces
- Monitor your health daily for symptoms

If you have any questions about biosecurity for horses, please contact your local extension agent. Happy riding!!

Virtual Livestock Options for NC Youth

By: Jamie Warner, Livestock Extension Agent with N.C. Cooperative Extension in Montgomery County

As if closing down the school system, churches, gyms, restaurants and more weren't enough, hundreds of youth livestock showmen across the state (and the nation, for that matter) were even more disappointed to learn that many of their favorite shows would also be canceled this year. In these times of uncertainty, these young people look to Showing as their "constant" and now that's not the case.

Enter NC Cooperative Extension. . .

NC Cooperative Extension Agriculture Agents realized that these young showmen had already had so much taken away from them this year that they did not want to see anything else be canceled, **IF** they could help it. Agents, in cooperation with the NC State Youth Livestock staff, put their heads together and transformed many of the planned events into Virtual Events that let youth participate, all the while maintaining social distance.

Programs like "How to Photograph Your Livestock", "Virtual Livestock Skillathons", "Virtual Livestock Judging" and more transformed into online platforms in order to allow young showmen to participate without the fear of getting exposed to COVID.

Showmanship Circuits were also impacted by the pandemic. Agriculture Agents from the South Central, North Central and Southeast Extension Districts worked together to turn the annual 4-H Farm Credit Showmanship Circuit into a "cyber circuit".

In July, the Circuit opened registration online and had over 70 showmen register to compete. Participants were given their first video prompt on August 1st and were given two weeks to shoot their videos. Videos were submitted on two weeks later to their County Ag Agents and judges watched and evaluated the videos before placing them based on showmen's knowledge of the prompt topic, animal handling and overall showmanship qualities. For prompt one, the showmen were tasked with introducing themselves, their animal and giving reasons why they chose that particular animal to show. They ended their videos by leading the animal around, setting the animals up and maintaining eye contact with the "judges" (through the camera). The second prompt was released to showmen on August 22nd and were submitted on August 29th. There will be 3 more prompts issued for a total of five. Participants are encouraged to complete all five but are only required to complete three in order to finish the circuit. Videos of the week for each species (cattle, goat and sheep) can be viewed on the Circuit Webpage at (<https://randolph.ces.ncsu.edu/videos-of-the-week/>).

Even though showing virtually isn't the preferred method of hands-on showing and learning, this virus epidemic has forced us all to think outside of the box, pulled us out of our comfort zones and made us rethink program presentations. It has compelled us to "Make The Best Better" (the 4-H Motto). To all showmen, we wish you luck and as always, if you need help, please contact your local Cooperative Extension Office. We are here to help you!



Water Quality

By: Richard Goforth, South Central Area Specialized Poultry Agent with N.C. Cooperative Extension

When is the last time you checked the quality of water your birds drink? While it is common for producers to check their water when they dig a new well or construct houses for many this is the last time they think about the source quality of their water. Most growers know they need to flush and clean their drinker lines regularly even if they don't admit they sometimes forget those steps. Many also have injection systems that strive to maintain the needed level of sanitizing agents to prevent bacteria growth in those drinker lines, but few really know the quality of the water they start with. Understanding what is in your water is key to managing its use and delivery and to understand water quality first it is important to know it can change. If water quality changes then it must be monitored to make sure it falls within the ranges for optimum bird performance. Many factors can affect water quality especially when groundwater is the primary source; which, is often the case for most poultry producers. Droughts, floods, and construction can all have major impacts on water quality and quantity. Water is the universal solvent so each time the water table changes the potential for new or greater levels of contamination to the ground water reservoir increase. New construction changes water runoff patterns, increases impervious surfaces and provides potential direct access to the reservoir with each new well. Even if you have municipal water there can be changes in quality that may affect the performance of sanitizers and vitamins; as well as, taste and bird acceptance. These often occur when switches are made in sanitizing agents or intake sources are changed.

Determining water quality starts with sampling your water, for both dissolved minerals and bacteria. Your county health department can usually provide bacteria testing for well water. Contact your county for scheduling, cost and procedures; many counties require a trained technician come out to take the sample to ensure there is no cross contamination. The presence of bacteria in water at the well can indicate issues with the well casing and often requires treatment to clear. Retesting will be required to assure the treatment worked and a follow up test within the year would be advised. An inspection of the lining should be considered especially if the treatment fails to reduce the load to an acceptable level or there is recurrence of the bacterial load. Testing for dissolved minerals in water is pretty simple in North Carolina as the NC Department of Agriculture & Consumer Services offers the standard test for only five dollars. To submit a sample all you have to do is remove the label from an empty plastic soda or water bottle and rinse it several times, do not use soap, from the water source you want to test, then fill it to the top leaving no air space and cap. Be sure to label the bottle with the same descriptive code you use on the submission form and send it with the fee to the lab promptly or refrigerate if it needs to be stored before submission. You can download the form and get more tips on sampling at <https://www.ncagr.gov/agronomi/uyrsoln.htm> .

Be sure to select the appropriate use code on the form to receive recommendations about acceptable levels of minerals in your water for poultry. Contact your Area Specialized Poultry agent, flock supervisor or water specialist for additional assistance in understanding your analysis and possible mediation needs. Remember that water quality not only affects bird performance it also has a big impact on equipment. Drinkers and cool cells may have different tolerances for contaminants than your birds. Consult your equipment manufacturer for recommendations and strategies to maintain equipment performance. To learn more read "Do you know your water Quality" at www.poultryventilation.com.